

Effect of the vaccination against Shiga toxin 2e in a farm with history of oedema disease, caused by atypical *Escherichia coli* producing Shiga toxin (STEC)

DANIEL SPERLING^{1*}, NAOMI ISAKA¹, HAMADI KAREMBE¹,
JONAS VANHARA^{1,4}, JOSEF VINDUSKA², NICOL STRAKOVA³,
ALZBETA KALOVA^{3,5}, IVANA KOLACKOVA³, RENATA KARPISKOVA³

¹*Ceva Sante Animale France, Libourne, France*

²*ZOD Zichlinek, Zichlinek, Czech Republic*

³*Veterinary Research Institute, Brno, Czech Republic*

⁴*Swine and Ruminant Clinic, Faculty of Veterinary Medicine, University of Veterinary Sciences Brno, Brno, Czech Republic*

⁵*Department of Experimental Biology, Faculty of Science, Masaryk University Brno, Brno, Czech Republic*

*Corresponding author: daniel.sperling@ceva.com

The authors are fully responsible for both the content and the formal aspects of the electronic supplementary material. No editorial adjustments were made.

Electronic Supplementary Material (ESM)

Figure S1. Flow chart – Vaccination against oedema disease (OD)




Figure S1. Flow chart – Vaccination against oedema disease (OD)

MMA = mastitis, metritis and agalactia syndrome in sows