

Ribavirin inhibits peste des petits ruminants virus proliferation *in vitro*

WEIFENG ZHANG¹, HUALONG DENG¹, YANFEN LIU¹, SHAOHONG CHEN²,
YOU LIU^{2*}, YUNTAO ZHAO^{2*}

¹Department of Animal Science, College of Coastal Agricultural Science,
Guangdong Ocean University, Zhanjiang, P.R. China

²Department of Bioengineering, College of Food Science and Technology,
Guangdong Ocean University, Zhanjiang, P.R. China

*Corresponding authors: liuy6254282@163.com; yuntaozhao@163.com

Weifeng Zhang and Hualong Deng contributed equally to this work

The authors are fully responsible for both the content and the formal aspects of the electronic supplementary material. No editorial adjustments were made.

Electronic Supplementary Material (ESM)

Figure S1. The impact of ribavirin on the expression levels of the essential proteins of the PI3K/AKT signalling pathway




Figure S1. The impact of ribavirin on the expression levels of the essential proteins of the PI3K/AKT signalling pathway. Vero cells were treated with ribavirin and infected with PPRV, and the expression levels of AKT, p-AKT, GSK3, p-GSK3, CREB, p-CREB, Bad, and p-Bad (A–H) were identified by western blotting 48 hours later. The data, obtained from three independent experiments, are presented as the mean \pm SD (&& P < 0.01, &&& P < 0.001 vs the virus control; * P < 0.05, ** P < 0.01, *** P < 0.001 vs the cell control)