https://doi.org/10.17221/76/2025-VETMED

Salmonella enterica isolates from poultry in North-Central Nigeria reveal multidrug-resistant patterns and risk of contamination to the human food chain

Abdullahi Ozomata Sanni[1,2], *Abdurrahman Hassan Jibril*[3], *Annelize Jonker*[1], *Folorunso Oludayo Fasina*[1,4]*

[1]*Department of Veterinary Tropical Diseases, University of Pretoria, Onderstepoort, South Africa*
[2]*Agro-Processing, Productivity Enhancement and Livelihood Improvement Support (APPEALS) Project, Lokoja, Kogi State, Nigeria*
[3]*Department of Veterinary Public Health and Preventive Medicine, Faculty of Veterinary Medicine, Usmanu Danfodiyo University, Sokoto, Sokoto State, Nigeria*
[4]*Department of Agriculture and Animal Health, College of Agriculture and Environmental Sciences, University of South Africa (UNISA), Florida Park, Roodepoort, South Africa*

**Corresponding author: daydupe2003@yahoo.co.uk*

The authors are fully responsible for both the content and the formal aspects of the electronic supplementary material. No editorial adjustments were made.

Electronic Supplementary Material (ESM)

Figure S1. Amplicons of *Salmonella* spp. visualised on stained 1.5% agarose gels

https://doi.org/10.17221/76/2025-VETMED




Figure S1. Amplicons of *Salmonella* spp. visualised on stained 1.5% agarose gels
The amplicon size was 284 bp measured against a 100 bp standard DNA ladder. The reference *Salmonella* strain (ATCC 14028) was used as positive control and sterile water without DNA was used as negative control
Note that only 42 *Salmonella* isolates were identified by PCR in this work