

***Salmonella enterica* isolates from poultry in North-Central Nigeria reveal multidrug-resistant patterns and risk of contamination to the human food chain**

ABDULLAHI OZOMATA SANNI^{1,2} , ABDURRAHMAN HASSAN JIBRIL³ ,
ANNELIZE JONKER¹ , FOLORUNSO OLUDAYO FASINA^{1,4*} 

¹Department of Veterinary Tropical Diseases, University of Pretoria, Onderstepoort, South Africa

²Agro-Processing, Productivity Enhancement and Livelihood Improvement Support (APPEALS) Project, Lokoja, Kogi State, Nigeria

³Department of Veterinary Public Health and Preventive Medicine, Faculty of Veterinary Medicine, Usmanu Danfodiyo University, Sokoto, Sokoto State, Nigeria

⁴Department of Agriculture and Animal Health, College of Agriculture and Environmental Sciences, University of South Africa (UNISA), Florida Park, Roodepoort, South Africa

*Corresponding author: daydupe2003@yahoo.co.uk

The authors are fully responsible for both the content and the formal aspects of the electronic supplementary material. No editorial adjustments were made.

Electronic Supplementary Material (ESM)

Figure S1. Amplicons of *Salmonella* spp. visualised on stained 1.5% agarose gels




Figure S1. Amplicons of *Salmonella* spp. visualised on stained 1.5% agarose gels

The amplicon size was 284 bp measured against a 100 bp standard DNA ladder. The reference *Salmonella* strain (ATCC 14028) was used as positive control and sterile water without DNA was used as negative control

Note that only 42 *Salmonella* isolates were identified by PCR in this work