

<https://doi.org/10.17221/56/2025-VETMED>

# Dairy bull sperm subpopulation behaviour in frozen-thawed semen across breed, temperature, and thawing time

FRANCISCO SEVILLA<sup>1,2</sup>, IGNACIO ARAYA-ZUNIGA<sup>2</sup>, JUAN MANUEL SOLIS<sup>2</sup>,  
CARINE CORCINI<sup>3</sup>, PATRICIA CERVANTES-ACOSTA<sup>4</sup>, ANTONIO HERNANDEZ-  
BELTRAN<sup>4</sup>, RAFAEL MOLINA-MONTERO<sup>5</sup>, DERLING PICHARDO-MATAMOROS<sup>6</sup>,  
ANTHONY VALVERDE<sup>2\*</sup>

<sup>1</sup>Doctorado en Ciencias Naturales para el Desarrollo (DOCINADE), Instituto Tecnológico de Costa Rica, Universidad Nacional, Universidad Estatal a Distancia, Alajuela, Costa Rica

<sup>2</sup>Laboratory of Animal Reproduction, School of Agronomy, Costa Rica Institute of Technology, Alajuela, Costa Rica

<sup>3</sup>Veterinary School, Pelotas Federal University, Rio Grande do Sul, Brazil

<sup>4</sup>School of Veterinary Medicine and Animal Husbandry, Veracruz University, Veracruz, México

<sup>5</sup>Agricultural Production Program, School of Agronomy, Costa Rica Institute of Technology, Alajuela, Costa Rica

<sup>6</sup>Animal Husbandry Engineering, National Agrarian University, Chontales, Nicaragua

\*Corresponding author: [anvalverde@itcr.ac.cr](mailto:anvalverde@itcr.ac.cr)

**Citation:** Sevilla F, Araya-Zuniga I, Solis JM, Corcini C, Cervantes-Acosta P, Hernandez-Beltran A, Molina-Montero R, Pichardo-Matamoros D, Valverde A (2026): Dairy bull sperm subpopulation behaviour in frozen-thawed semen across breed, temperature, and thawing time. *Vet Med-Czech* 71, 1–9.

**Abstract:** Optimising dairy cattle reproduction with assisted reproductive technologies, such as artificial insemination, requires standardised semen handling and analysis. This study evaluated the kinematic structure of the spermatozoan subpopulation in frozen-thawed Holstein and Jersey bull semen doses under different thawing protocols. We used frozen semen doses taken from four bulls of each breed. Nine semen doses were collected from each animal, for a total of 72 cryopreserved doses. Straw thawing was performed at three temperatures (35 °C, 37 °C, and 40 °C) and three times (30 s, 40 s, and 45 s). Sperm kinematic patterns were evaluated using a CASA-mot system (ISAS<sup>®</sup>v1). Sperm kinematic variables for each breed, temperature, and thawing time identified four subpopulations. The analysis revealed an effect of breed and semen thawing protocols ( $P < 0.05$ ) on sperm subpopulation distribution, sperm movement, and swimming patterns. Subpopulation analyses based on semen assessment are needed to further interpret the relevance and effect on fertility.

**Keywords:** cryopreservation; dairy cattle; reproduction; spermatozoa; sperm kinematics

Funded by the Costa Rica Institute of Technology Vice-Chancellor's Office of Research and Extension. This work was part of the research project VIE-2151083, Optimización de la conservación y búsqueda de parámetros de la fertilidad en espermatozoides de animales de interés productivo. The Article Processing Charges were covered by the Costa Rica Institute of Technology.

© The authors. This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International (CC BY-NC 4.0).

Animal reproduction is central to dairy production performance (Cheng et al. 2022). Assisted reproduction techniques, such as artificial insemination with fresh or frozen-thawed semen, improve animal genetics (Souza et al. 2016; Koch et al. 2022). Success of artificial insemination, however, depends on environmental conditions, management, nutrition, and semen quality (Barquero et al. 2021a; Koch et al. 2022). These reproductive biotechnologies require clear, validated, replicable protocols that minimise external variation in livestock systems (Mikkola et al. 2024). Studies have shown that protocols that manipulate semen during seminal sample thawing can influence sperm motility and functionality. Such variables include percentage of total and progressive sperm motility, plasma membrane integrity, and sperm kinetics (Diskin 2018; Goshme et al. 2021; Solis et al. 2024).

Semen analysis has traditionally focused on cell motility or morphology by studying the percentage of sperm with normal morphology in an ejaculate or seminal dose (Silvestre et al. 2023). With the emergence of Computer-Assisted Semen Analysis (CASA) systems that can now generate larger volumes of data with high-resolution cameras (Bompart et al. 2019; Valverde et al. 2019), sperm movement patterns can be predicted with greater accuracy and precision using multivariate statistical analysis models. This kind of analysis can better determine semen quality and predict male fertility (Martinez-Pastor et al. 2011; Hidalgo et al. 2021; Martinez-Pastor 2022). The analysis of cellular subsets within the ejaculate using multivariate data analysis has, furthermore, allowed us to identify sperm subpopulations within an ejaculate (Valverde et al. 2016; Martinez-Pastor 2022). These subpopulations exhibit different movement patterns that reorganise themselves based on intrinsic and extrinsic factors (Murphy et al. 2018) indicative of different sperm functions (Martinez et al. 2013; Santolaria et al. 2015; Ibanescu et al. 2018; Azevedo et al. 2024).

Sperm subpopulations of various ruminants have been classified based on motility and kinematic characteristics (Perez-Marin et al. 2021; Kanno et al. 2023), morphometry (Maroto-Morales et al. 2012; Valverde et al. 2016; Viquez et al. 2023), as well as other external factors (Garcia-Herreros and Leal 2014) such as sexual status (Araya-Zuniga et al. 2024), ejaculate fractions (Perez-Marin et al.

2021), species (Gacem et al. 2021), season (Ibanescu et al. 2018) and age (Marti et al. 2011; Menon et al. 2011). With subpopulation classification, we can now analyse sperm distribution within the ejaculate and identify animals that harbour sperm subpopulations with distributions that favour reproductive fitness and suitability for breeding or artificial insemination (Hidalgo et al. 2022). This study evaluated the kinematic behaviour of sperm subpopulations in frozen-thawed seminal doses of Holstein and Jersey bulls with different thawing protocols.

## MATERIAL AND METHODS

### Ethical approval

The study was conducted in compliance with Costa Rica's national regulations governing research with live animals. All procedures were performed with care to minimise stress and ensure animal welfare. Ethical approval was granted by the Committee of the Tropical Sustainable Agriculture Research and Development Center (CIDASTH-ITCR), in accordance with Section 08/2023 and Article 5.0 of DAGSC-075-2023 and CIE-206-2023. The study also complied with the ARRIVE guidelines (<https://arriveguidelines.org/>) for reporting standards in animal research.

### Study location, seminal doses, and thawing protocols

This study was conducted at the Costa Rica Institute of Technology's Animal Reproduction Laboratory at the San Carlos Campus in Alajuela, Costa Rica, from April to November 2023. Semen dose quality was verified prior to cryopreservation, and only ejaculates with sperm motility above 60% and normal morphology above 75% were used. Nine cryopreserved seminal doses were collected from eight bulls (Holstein,  $n = 4$ ; Jersey,  $n = 4$ ), totalling 72 doses. These doses were packaged in 0.25 ml volume straws and stored in an MVE<sup>®</sup> XC 20 Signature cryogenic tank (MVE Biological Solutions, Ball Ground, GA, USA). To thaw doses, we followed the methodology described in Solis et al. (2024). To thaw straws in a thermoregulated water bath (Digisystem Laboratory Instruments

<https://doi.org/10.17221/56/2025-VETMED>

Inc., New Taipei City, Taiwan), three temperatures (35 °C, 37 °C, and 40 °C) and three times (30 s, 40 s and 45 s) were applied. After thawing, the contents of each straw were placed in an Eppendorf® tube (Sigma-Aldrich, St. Louis, MO, USA). The samples were diluted with the commercial diluent OptiXcell® (IMV, L'Aigle, France) in a 1 : 1 (vol : vol) ratio, for an average final concentration of  $33.19 \pm 13.46 \times 10^6/\text{ml}$  per straw for the Jersey breed and  $21.61 \pm 14.54 \times 10^6/\text{ml}$  per straw for the Holstein.

### Sperm kinematic assessment

To assess kinematics, a previously preheated at 37 °C sample was homogenised by gentle shaking, and then a 3 µl diluted semen subsample was extracted and placed in a Spermtrack® sperm counting chamber (Proiser I+D, Paterna, Spain) with a depth of 20 µm. The counting chamber had been preheated to 37 °C on a hot plate.

A CASA-mot ISAS®v1 system (Integrated Semen Analysis System, Proiser I+D, Paterna, Spain) was used. The system consisted of a video camera (Proiser 782M, Proiser I+D) and a UB203 microscope (UOP/Proiser R+D) equipped with a 1× eyepiece and a 10× negative phase-contrast objective (NA 0.25). The camera recorded at 50 fps with a final resolution of  $768 \times 576$  pixels. The sperm counting chamber containing the sample was placed on the microscope's integrated, preheated fixed stage at  $37.0 \pm 0.5$  °C.

Two replicates of each sample were analysed, and a total of seven microscopic fields or a minimum of 600 spermatozoa were captured for each. Kinematic analysis included rectilinear velocity (VSL,  $\mu\text{m s}^{-1}$ ), curvilinear velocity (VCL,  $\mu\text{m s}^{-1}$ ), average path velocity (VAP,  $\mu\text{m s}^{-1}$ ), and percentages of linearity ( $\text{LIN} = \text{VSL}/\text{VCL} \times 100$ ), straightness ( $\text{STR} = \text{VSL}/\text{VAP} \times 100$ ), and wobble ( $\text{WOB} = \text{VAP}/\text{VCL} \times 100$ ). The crossover frequency (BCF, in Hz) and lateral head displacement (ALH, in µm) were also analysed.

### Statistical analysis

Results were presented as mean  $\pm$  standard error of the mean. Assumptions of normality and homoscedasticity were assessed using the Shapiro–

Wilk–Levene test. Once these assumptions were verified, a repeated-means analysis of variance (ANOVA) was performed. General linear and mixed models were used to analyse sperm kinematic variables. Means were compared using the least-squares method with Bonferroni correction at  $P < 0.05$ . Kinematic variables (VCL, VAP, VSL, LIN, STR, WOB, ALH, and BCF) were divided into subsets by breed, temperature, and thawing time, and multivariate procedures then identified groups within each subset. Variables were standardised, followed by a principal factor analysis (PFA). PFs with an eigenvalue greater than one were extracted from the PFA using the Kaiser criterion, and the Kaiser–Meyer–Olkin (KMO) test was applied. PFs were obtained by applying the Varimax rotation method with Kaiser normalisation. A non-hierarchical K-means cluster analysis was used to identify sperm subpopulations, and ANOVA was used to evaluate statistical differences between groups obtained for all kinematic variables. A post hoc pairwise comparison of group means was performed using the Tukey–Kramer test, with  $P < 0.05$ . All analyses were performed using IBM SPSS v23.0 for Windows (SPSS Inc., Chicago, IL, USA).

## RESULTS

PFA identified four sperm subpopulations in each breed. The Holstein subpopulation 4 (SP<sub>4</sub>) had the highest VCL, while SP<sub>1</sub> had sperm with the highest VSL and VAP. The SPs with the most linear movements, as described by LIN, STR, and WOB, were SP<sub>1</sub> and SP<sub>3</sub>. In the Jersey breed, the highest VCL was observed in SP<sub>4</sub>. The highest averages for VSL and VAP were found in SP<sub>1</sub>. There was variation in progressivity because the most linear and oscillating movement was exhibited by SP<sub>3</sub>, while the greatest straightness was exhibited by SP<sub>1</sub> (Table 1).

Significant differences in ALH were found among subpopulations by breed ( $P < 0.05$ ). Mean values in SP<sub>4</sub> for both breeds were higher than in other subpopulations ( $P < 0.05$ ). SP<sub>3</sub> for both Holstein and Jersey had the lowest means among subpopulations. BCF showed statistically significant differences for subpopulations within each breed ( $P < 0.05$ ). The highest BCF values were observed in SP<sub>1</sub> for both, while the lowest were observed in SP<sub>2</sub> (Figure 1).

Table 1. Subpopulations (mean  $\pm$  SEM) according to frozen-thawed semen kinematic variables by dairy cattle breed

Variable	VCL	VSL	VAP	LIN	STR	WOB
Holstein						
SP <sub>1</sub>	190.46 $\pm$ 0.28 <sup>b</sup>	113.86 $\pm$ 0.22 <sup>a</sup>	118.69 $\pm$ 0.20 <sup>a</sup>	59.52 $\pm$ 0.10 <sup>b</sup>	93.97 $\pm$ 0.16 <sup>a</sup>	62.27 $\pm$ 0.09 <sup>b</sup>
SP <sub>2</sub>	97.94 $\pm$ 0.21 <sup>d</sup>	25.73 $\pm$ 0.16 <sup>d</sup>	44.67 $\pm$ 0.15 <sup>d</sup>	26.11 $\pm$ 0.08 <sup>d</sup>	57.41 $\pm$ 0.12 <sup>d</sup>	45.95 $\pm$ 0.07 <sup>c</sup>
SP <sub>3</sub>	99.13 $\pm$ 0.29 <sup>c</sup>	67.19 $\pm$ 0.22 <sup>c</sup>	70.81 $\pm$ 0.21 <sup>c</sup>	66.74 $\pm$ 0.11 <sup>a</sup>	91.23 $\pm$ 0.17 <sup>b</sup>	71.36 $\pm$ 0.10 <sup>a</sup>
SP <sub>4</sub>	213.02 $\pm$ 0.36 <sup>a</sup>	70.08 $\pm$ 0.28 <sup>b</sup>	95.52 $\pm$ 0.26 <sup>b</sup>	32.50 $\pm$ 0.13 <sup>c</sup>	71.86 $\pm$ 0.21 <sup>c</sup>	44.90 $\pm$ 0.12 <sup>d</sup>
Jersey						
SP <sub>1</sub>	209.52 $\pm$ 0.28 <sup>b</sup>	131.74 $\pm$ 0.22 <sup>a</sup>	137.69 $\pm$ 0.20 <sup>a</sup>	62.58 $\pm$ 0.09 <sup>b</sup>	93.40 $\pm$ 0.13 <sup>a</sup>	65.76 $\pm$ 0.08 <sup>b</sup>
SP <sub>2</sub>	100.67 $\pm$ 0.26 <sup>c</sup>	26.30 $\pm$ 0.21 <sup>d</sup>	47.29 $\pm$ 0.19 <sup>d</sup>	25.88 $\pm$ 0.09 <sup>d</sup>	55.01 $\pm$ 0.12 <sup>d</sup>	47.19 $\pm$ 0.07 <sup>c</sup>
SP <sub>3</sub>	110.91 $\pm$ 0.28 <sup>d</sup>	79.36 $\pm$ 0.23 <sup>c</sup>	84.06 $\pm$ 0.20 <sup>c</sup>	70.21 $\pm$ 0.09 <sup>a</sup>	91.28 $\pm$ 0.13 <sup>b</sup>	75.22 $\pm$ 0.08 <sup>a</sup>
SP <sub>4</sub>	250.35 $\pm$ 0.33 <sup>a</sup>	88.90 $\pm$ 0.27 <sup>b</sup>	116.54 $\pm$ 0.24 <sup>b</sup>	34.88 $\pm$ 0.11 <sup>c</sup>	74.13 $\pm$ 0.15 <sup>c</sup>	46.72 $\pm$ 0.09 <sup>d</sup>

<sup>a-d</sup>Different letters between columns indicate differences between subpopulations according to breed ( $P < 0.05$ )

LIN = linearity index (%); SEM = standard error of the mean; SP = subpopulation; STR = straightness index (%); VAP = average path velocity ( $\mu\text{m s}^{-1}$ ); VCL = curvilinear velocity ( $\mu\text{m s}^{-1}$ ); VSL = rectilinear velocity ( $\mu\text{m s}^{-1}$ ); WOB = trajectory oscillation (%)




Figure 1. Mean oscillation values by sperm subpopulation in frozen-thawed dairy cattle semen samples

<sup>a-d</sup>Different letters show differences between subpopulations within each breed ( $P < 0.05$ )

ALH ( $\mu\text{m}$ ) = amplitude of lateral head displacement; BCF (Hz) = beat-cross frequency

Subpopulations were characterised according to semen thawing temperatures. At 35 °C, differences were observed between sperm subpopulations ( $P < 0.05$ ), with the highest values in SP<sub>4</sub>, followed by SP<sub>1</sub>, SP<sub>3</sub>, and SP<sub>2</sub>. This pattern was similar for semen thawing temperatures of 37 °C and 40 °C. Overall, SP<sub>3</sub> was the subpopulation that showed the most progressive movement across all temperatures (Table 2).

At a thawing temperature of 35 °C, the greatest sperm undulation was observed in SP<sub>4</sub> and SP<sub>1</sub>, with higher average ALH and BCF values, respectively. A similar pattern was observed at temperatures of 37 °C and 40 °C (Figure 2). SP<sub>2</sub> and SP<sub>3</sub> had

the lowest average values across all analysed temperatures ( $P < 0.05$ ).

The sperm subpopulation was analysed for thawing times (Table 3). For the 30 s thawing time, significant differences were observed between subpopulations ( $P < 0.05$ ). The SP<sub>4</sub> subpopulation had the highest velocity, while the cells with the most progressive, linear movement were present in SP<sub>3</sub>. At the thawing times of 40 s and 45 s, very similar subpopulation behaviour patterns emerged, with the highest velocity and progressiveness observed in SP<sub>1</sub> and SP<sub>3</sub>, respectively. There were differences between subpopulations for the three thawing temperatures ( $P < 0.05$ ) in the sperm oscillation

<https://doi.org/10.17221/56/2025-VETMED>

Table 2. Subpopulations (mean ± SEM) according to kinematic variables of frozen-thawed semen and thawing temperature

Variable	VCL	VSL	VAP	LIN	STR	WOB
35 °C						
SP <sub>1</sub>	202.21 ± 0.37 <sup>b</sup>	125.80 ± 0.29 <sup>a</sup>	130.86 ± 0.27 <sup>a</sup>	61.87 ± 0.12 <sup>b</sup>	93.81 ± 0.18 <sup>a</sup>	64.70 ± 0.11 <sup>b</sup>
SP <sub>2</sub>	97.75 ± 0.31 <sup>d</sup>	25.27 ± 0.24 <sup>d</sup>	45.21 ± 0.22 <sup>d</sup>	25.72 ± 0.10 <sup>d</sup>	55.71 ± 0.15 <sup>d</sup>	46.69 ± 0.09 <sup>c</sup>
SP <sub>3</sub>	104.16 ± 0.38 <sup>c</sup>	73.52 ± 0.30 <sup>c</sup>	77.82 ± 0.27 <sup>c</sup>	68.97 ± 0.13 <sup>a</sup>	90.82 ± 0.19 <sup>b</sup>	74.13 ± 0.11 <sup>a</sup>
SP <sub>4</sub>	236.29 ± 0.45 <sup>a</sup>	81.87 ± 0.35 <sup>b</sup>	107.98 ± 0.33 <sup>b</sup>	33.82 ± 0.15 <sup>c</sup>	73.27 ± 0.22 <sup>c</sup>	45.74 ± 0.13 <sup>d</sup>
37 °C						
SP <sub>1</sub>	202.13 ± 0.34 <sup>b</sup>	122.94 ± 0.26 <sup>a</sup>	128.69 ± 0.24 <sup>a</sup>	60.47 ± 0.11 <sup>b</sup>	93.53 ± 0.16 <sup>a</sup>	63.58 ± 0.10 <sup>b</sup>
SP <sub>2</sub>	100.66 ± 0.29 <sup>d</sup>	26.75 ± 0.22 <sup>d</sup>	46.45 ± 0.21 <sup>d</sup>	26.36 ± 0.10 <sup>d</sup>	57.17 ± 0.14 <sup>d</sup>	46.29 ± 0.08 <sup>c</sup>
SP <sub>3</sub>	107.80 ± 0.36 <sup>c</sup>	74.88 ± 0.28 <sup>c</sup>	79.05 ± 0.26 <sup>c</sup>	68.30 ± 0.12 <sup>a</sup>	91.59 ± 0.17 <sup>b</sup>	72.89 ± 0.10 <sup>a</sup>
SP <sub>4</sub>	236.12 ± 0.41 <sup>a</sup>	82.28 ± 0.31 <sup>b</sup>	109.23 ± 0.29 <sup>b</sup>	34.23 ± 0.14 <sup>c</sup>	73.41 ± 0.20 <sup>c</sup>	46.30 ± 0.12 <sup>c</sup>
40 °C						
SP <sub>1</sub>	200.19 ± 0.36 <sup>b</sup>	124.27 ± 0.29 <sup>a</sup>	129.85 ± 0.27 <sup>a</sup>	61.71 ± 0.13 <sup>b</sup>	93.57 ± 0.18 <sup>a</sup>	64.75 ± 0.11 <sup>b</sup>
SP <sub>2</sub>	99.18 ± 0.30 <sup>d</sup>	25.94 ± 0.24 <sup>d</sup>	46.04 ± 0.22 <sup>d</sup>	25.89 ± 0.11 <sup>d</sup>	55.84 ± 0.15 <sup>d</sup>	46.66 ± 0.09 <sup>c</sup>
SP <sub>3</sub>	106.10 ± 0.36 <sup>c</sup>	74.63 ± 0.28 <sup>c</sup>	78.92 ± 0.26 <sup>c</sup>	69.09 ± 0.12 <sup>a</sup>	91.32 ± 0.18 <sup>b</sup>	73.93 ± 0.11 <sup>a</sup>
SP <sub>4</sub>	234.70 ± 0.46 <sup>a</sup>	80.40 ± 0.36 <sup>b</sup>	107.52 ± 0.34 <sup>b</sup>	33.73 ± 0.16 <sup>c</sup>	72.99 ± 0.23 <sup>c</sup>	45.90 ± 0.14 <sup>d</sup>

<sup>a-d</sup>Different letters between columns indicate differences between subpopulations according to semen thawing temperature ( $P < 0.05$ )

LIN = linearity index (%); SEM = standard error of the mean; SP = subpopulation; STR = straightness index (%); VAP = average path velocity ( $\mu\text{m s}^{-1}$ ); VCL = curvilinear velocity ( $\mu\text{m s}^{-1}$ ); VSL = rectilinear velocity ( $\mu\text{m s}^{-1}$ ); WOB = trajectory oscillation (%)




Figure 2. Mean sperm oscillation values obtained from frozen-thawed semen samples from dairy cattle at different thawing temperatures

<sup>a-d</sup>Different letters show differences between subpopulations within each thawing temperature ( $P < 0.05$ )

ALH (µm) = amplitude of lateral head displacement; BCF (Hz) = beat-cross frequency

variables, ALH and BCF; however, no differences were found ( $P > 0.05$ ) between semen thawing temperatures within subpopulations. The highest ALH

values were observed in SP<sub>4</sub>, with values exceeding 5 µm. For BCF, the highest averages were in SP<sub>4</sub> across all thawing temperatures (Figure 3).

Table 3. Kinematic subpopulations of frozen-thawed semen from dairy cattle according to the thawing time

Variable	VCL	VSL	VAP	LIN	STR	WOB
30 s						
SP <sub>1</sub>	201.18 ± 0.35 <sup>b</sup>	123.38 ± 0.27 <sup>b</sup>	128.96 ± 0.25 <sup>a</sup>	60.95 ± 0.12 <sup>b</sup>	93.57 ± 0.17 <sup>a</sup>	64.01 ± 0.10 <sup>b</sup>
SP <sub>2</sub>	100.03 ± 0.29 <sup>d</sup>	26.19 ± 0.23 <sup>d</sup>	46.20 ± 0.21 <sup>d</sup>	25.94 ± 0.10 <sup>d</sup>	56.17 ± 0.14 <sup>d</sup>	46.42 ± 0.08 <sup>c</sup>
SP <sub>3</sub>	106.66 ± 0.36 <sup>c</sup>	74.36 ± 0.28 <sup>c</sup>	78.59 ± 0.26 <sup>c</sup>	68.60 ± 0.12 <sup>a</sup>	91.33 ± 0.18 <sup>b</sup>	73.37 ± 0.10 <sup>a</sup>
SP <sub>4</sub>	236.29 ± 0.42 <sup>a</sup>	81.70 ± 0.32 <sup>a</sup>	108.33 ± 0.30 <sup>b</sup>	33.97 ± 0.14 <sup>c</sup>	73.37 ± 0.20 <sup>c</sup>	45.93 ± 0.12 <sup>d</sup>
40 s						
SP <sub>1</sub>	201.64 ± 0.36 <sup>b</sup>	124.84 ± 0.29 <sup>a</sup>	130.24 ± 0.26 <sup>a</sup>	61.60 ± 0.12 <sup>b</sup>	93.37 ± 0.17 <sup>a</sup>	64.56 ± 0.11 <sup>b</sup>
SP <sub>2</sub>	99.04 ± 0.32 <sup>d</sup>	26.18 ± 0.25 <sup>d</sup>	45.74 ± 0.23 <sup>d</sup>	26.22 ± 0.11 <sup>d</sup>	56.88 ± 0.15 <sup>d</sup>	46.39 ± 0.09 <sup>c</sup>
SP <sub>3</sub>	107.71 ± 0.38 <sup>c</sup>	75.92 ± 0.30 <sup>c</sup>	80.31 ± 0.28 <sup>c</sup>	69.06 ± 0.13 <sup>a</sup>	91.38 ± 0.18 <sup>b</sup>	73.94 ± 0.11 <sup>a</sup>
SP <sub>4</sub>	235.58 ± 0.46 <sup>a</sup>	82.47 ± 0.36 <sup>b</sup>	108.73 ± 0.33 <sup>b</sup>	34.37 ± 0.16 <sup>c</sup>	73.85 ± 0.20 <sup>c</sup>	46.21 ± 0.13 <sup>c</sup>
45 s						
SP <sub>1</sub>	201.88 ± 0.36 <sup>b</sup>	124.66 ± 0.28 <sup>a</sup>	130.11 ± 0.26 <sup>a</sup>	61.39 ± 0.12 <sup>b</sup>	93.61 ± 0.18 <sup>a</sup>	64.35 ± 0.11 <sup>b</sup>
SP <sub>2</sub>	98.57 ± 0.29 <sup>d</sup>	25.66 ± 0.23 <sup>d</sup>	45.77 ± 0.21 <sup>d</sup>	25.86 ± 0.10 <sup>d</sup>	55.83 ± 0.15 <sup>d</sup>	46.80 ± 0.09 <sup>c</sup>
SP <sub>3</sub>	104.05 ± 0.36 <sup>c</sup>	73.00 ± 0.28 <sup>c</sup>	77.15 ± 0.26 <sup>c</sup>	68.73 ± 0.12 <sup>a</sup>	91.07 ± 0.18 <sup>b</sup>	73.64 ± 0.11 <sup>a</sup>
SP <sub>4</sub>	235.30 ± 0.44 <sup>a</sup>	80.50 ± 0.34 <sup>b</sup>	107.94 ± 0.32 <sup>b</sup>	33.54 ± 0.15 <sup>c</sup>	72.53 ± 0.22 <sup>c</sup>	45.91 ± 0.13 <sup>c</sup>

<sup>a-d</sup>Different letters between columns indicate differences between subpopulations according to the thawing time of seminal doses ( $P < 0.05$ )

LIN = linearity index (%); SEM = standard error of the mean; SP = subpopulation; STR = straightness index (%); VAP = average path velocity ( $\mu\text{m s}^{-1}$ ); VCL = curvilinear velocity ( $\mu\text{m s}^{-1}$ ); VSL = rectilinear velocity ( $\mu\text{m s}^{-1}$ ); WOB = trajectory oscillation (%)




Figure 3. Mean sperm oscillation values obtained from frozen-thawed dairy cattle semen at different thawing times

<sup>a-d</sup>Different letters show differences between subpopulations within each thawing time ( $P < 0.05$ )

ALH ( $\mu\text{m}$ ) = amplitude of lateral head displacement; BCF (Hz) = beat-cross frequency

## DISCUSSION

Seminal doses used in assisted reproduction processes exhibit significant variations in movement patterns and kinematics, as evidenced by groupings within each dose (Muino et al. 2008; Muino et al. 2009; Yaniz et al. 2018; Ibanescu et al. 2020). These

variations may be influenced by individual variables (e.g., breed) and ejaculate handling (temperature and thawing time) (Valverde et al. 2016). Variable sperm behaviour in *Bos taurus* bull ejaculates across breeds has already been reported (Menon et al. 2011; Ntemka et al. 2016). This variability contributes to its adaptability to the environmental

<https://doi.org/10.17221/56/2025-VETMED>

conditions in which individuals develop. The protocols used to thaw semen can provoke external variation in kinematic cell patterns (Alvarez-Rodriguez et al. 2024; Solis et al. 2024). The results of this work indicate that different semen temperatures and thawing times affect swimming patterns; furthermore, this may affect the probability of sperm fertilising once doses are used (Koch et al. 2022).

Sperm motility is a principal fertility indicator, since the sperm, once it enters the female reproductive tract, must traverse a long gauntlet of physical and chemical obstacles to reach the ampulla (Ntemka et al. 2016). Previous studies show that Holstein and Jersey breeds exhibit differences in sperm motility and kinematic patterns (Solis et al. 2024); however, subpopulation characterisation was not done. This underscores the utility of subpopulation dynamics analysis for interpreting the biological implications of ejaculate heterogeneity and its potential impact on fertility (Valverde et al. 2019; Azevedo et al. 2024; Martinez-Pastor 2022). These analyses elucidate male fertility by understanding sperm behaviour within the ejaculate and its potential functionality (Keyer et al. 2022). This knowledge provides a more objective criterion for selecting animals to participate in genetic improvement programs at artificial insemination centres (Ferraz et al. 2014).

Previous cluster analysis and identification of sperm subpopulations within ejaculate fractions (Valverde et al. 2016; Perez-Marin et al. 2021) have demonstrated ejaculate variability and the need to select among those fractions for the greatest preservation value. This discovery has recently led to numerous studies linking fertility to sperm subpopulation dynamics (Hidalgo et al. 2021; Hidalgo et al. 2022; Alm-Kristiansen 2023). It has further allowed us to associate swimming patterns, speed, and linearity with higher fertility rates (Ahmed et al. 2016; Yaniz et al. 2016; Pena et al. 2018). Other studies have shown that low-fertility males suffer from ejaculate subpopulations dominated by slow, non-linear movements (Ramon et al. 2013).

Some authors hypothesise that different subpopulations perform different functions within the ejaculate (Ibanescu et al. 2020). This functionality and group distribution vary depending on ejaculate-handling conditions and breed (Garcia-Alvarez et al. 2014; Alm-Kristiansen 2023). Overall, this study shows that kinematic differences are higher in two of the four identified subpopulations. These variations may

be due to the individual nature of the animal and its ejaculate per se (Fernandez-Lopez et al. 2022), as well as different functional subpopulations within the ejaculate and sperm maturity (Alm-Kristiansen 2023). Despite these advancements, it is still not possible to precisely determine the effects of these functional subpopulations on fertility (Barquero et al. 2021b). Our results also show that sperm distribution and kinematic patterns in subpopulations were more progressive and linear, depending on the thawing time and temperature protocol used, an observation not previously reported. This new evidence calls for additional study of these variables and their potential impact on sperm function, especially under tropical conditions.

In conclusion, the aim of this study was to investigate the effect of animal breed and semen thawing protocols on sperm subpopulation dynamics. Based on the data from this study, we observed that within the practical ranges tested (35–40 °C and 30–45 s), changing thawing temperature or thawing time does not eliminate or fundamentally alter the key progressive subpopulations (SP<sub>1</sub> and SP<sub>3</sub>). This suggests that the structure of post-thaw sperm motility and kinematics is robust to small protocol variations and that the functionally relevant progressive fraction is preserved. Subpopulation analysis is needed to improve the interpretation of its biological relevance and effect on fertility.

## Acknowledgement

The authors thank the Costa Rica Institute of Technology for funding this study. The funders had no role in the study design, data collection and analysis, decision to publish, or manuscript preparation.

## Conflict of interest

The authors declare no conflict of interest.

## REFERENCES

- Ahmed H, Andrabi SMH, Jahan S. Semen quality parameters as fertility predictors of water buffalo bull spermatozoa during low-breeding season. *Theriogenology*. 2016 Oct 1;86(6):1516-22.

<https://doi.org/10.17221/56/2025-VETMED>

- Alm-Kristiansen AH. Motility subpopulations with distinct motility characteristics using swim-up-selected sperm cells from Norwegian Red bulls: Effects of freezing-thawing and between-bull variation. *Biology (Basel)*. 2023 Aug 3;12(8):1086.
- Alvarez-Rodriguez M, Tomas-Almenar C, Nieto-Cristobal H, de Mercado E. Evaluation of different thawing protocols on Iberian boar sperm preserved for 10 years at different liquid nitrogen levels. *Animals (Basel)*. 2024 Mar 15;14(6):914.
- Araya-Zuniga I, Sevilla F, Molina-Montero R, Roldan ERS, Barrientos-Morales M, Silvestre MA, Valverde A. Kinematic and morphometric assessment of fresh semen, before, during and after mating period in Brahman bulls. *Animals (Basel)*. 2024 Jan 3;14(1):132.
- Azevedo HC, Blackburn HD, Lozada-Soto EA, Spiller SE, Purdy PH. Enhancing evaluation of bull fertility through multivariate analysis of sperm. *J Dairy Sci*. 2024 Dec; 107(12):11774-84.
- Barquero V, Roldan ERS, Soler C, Yaniz JL, Camacho M, Valverde A. Predictive capacity of boar sperm morphology and morphometric sub-populations on reproductive success after artificial insemination. *Animals (Basel)*. 2021a Mar 24;11(4):920.
- Barquero V, Roldan ERS, Soler C, Vargas-Leiton B, Sevilla F, Camacho M, Valverde A. Relationship between fertility traits and kinematics in clusters of boar ejaculates. *Biology (Basel)*. 2021b Jun 28;10(7):595.
- Bompard D, Vazquez RF, Gomez R, Valverde A, Roldan ERS, Garcia-Molina A, Soler C. Combined effects of type and depth of counting chamber, and rate of image frame capture, on bull sperm motility and kinematics. *Anim Reprod Sci*. 2019 Oct;209:106169.
- Cheng M, McCarl B, Fei C. Climate change and livestock production: A literature review. *Atmosphere*. 2022 Jan 15; 13(1):140.
- Diskin MG. Review: Semen handling, time of insemination and insemination technique in cattle. *Animal*. 2018 Jun; 12(s1):s75-84.
- Fernandez-Lopez P, Garriga J, Casas I, Yeste M, Bartumeus F. Predicting fertility from sperm motility landscapes. *Commun Biol*. 2022 Sep 28;5(1):1027.
- Ferraz MA, Morato R, Yeste M, Arcarons N, Pena AI, Tamarco C, Hidalgo CO, Muino R, Mogas T. Evaluation of sperm subpopulation structure in relation to in vitro sperm-oocyte interaction of frozen-thawed semen from Holstein bulls. *Theriogenology*. 2014 May;81(8):1067-72.
- Gacem S, Valverde A, Catalan J, Yanez Ortiz I, Soler C, Miro J. A new approach of sperm motility subpopulation structure in donkey and horse. *Front Vet Sci*. 2021 May 25;8:651477.
- Garcia-Alvarez O, Maroto-Morales A, Ramon M, del Olmo E, Jimenez-Rabadan P, Fernandez-Santos MR, Anel-Lopez L, Garde JJ, Soler AJ. Dynamics of sperm subpopulations based on motility and plasma membrane status in thawed ram spermatozoa incubated under conditions that support in vitro capacitation and fertilisation. *Reprod Fertil Dev*. 2014 Jun;26(5):725-32.
- Garcia-Herreros M, Leal CL. Comparative study of sperm washing and selection methods after cryopreservation and its influence on sperm subpopulational structure in a bovine model. *Syst Biol Reprod Med*. 2014 Dec;60(6): 338-47.
- Goshme S, Asfaw T, Demiss C, Besufekad S. Evaluation of motility and morphology of frozen bull semen under different thawing methods used for artificial insemination in North Shewa zone, Ethiopia. *Heliyon*. 2021 Oct 13; 7(10):e08183.
- Hidalgo MMT, Almeida ABM, Moraes FLZ, Marubayashi RYP, Souza FF, Barreiros TRR, Martins MIM. Sperm subpopulations influence the pregnancy rates in cattle. *Reprod Domest Anim*. 2021 Aug;56(8):1117-27.
- Hidalgo MMT, Marques de Almeida AB, Zito de Moraes FL, Marubayashi RYP, Palhaci RY, Barreiros TRR, Martins MIM. Influence of sperm post-thaw subpopulations of Angus and Nelore bulls on pregnancy rates by fixed-time artificial insemination. *Cryo Letters*. 2022 May-Jun;43(3):140-9.
- Ibanescu I, Leiding C, Bollwein H. Cluster analysis reveals seasonal variation of sperm subpopulations in extended boar semen. *J Reprod Dev*. 2018 Feb 27;64(1):33-9.
- Ibanescu I, Siuda M, Bollwein H. Motile sperm subpopulations in bull semen using different clustering approaches – Associations with flow cytometric sperm characteristics and fertility. *Anim Reprod Sci*. 2020 Apr;215:106329.
- Kanno C, Sakamoto KQ, Kang SS, Yanagawa Y, Katagiri S, Nagano M. The difference in subpopulation structures in sex-sorted and non-sorted semen by discriminant analysis of bull sperm motility data collected by a Computer-Assisted Sperm Analysis System. *Andrologia*. 2023; 2023:9944344.
- Keyer S, Van Der Horst G, Maree L. New approaches to define the functional competency of human sperm subpopulations and its relationship to semen quality. *Int J Fertil Steril*. 2022 Aug 21;16(3):140-51.
- Koch J, Weber LP, Heppelmann M, Freise F, Klingelmann M, Bachmann L. Effect of different thawing methods for frozen bull semen and additional factors on the conception rate of dairy cows in artificial insemination. *Animals (Basel)*. 2022 Sep 7;12(18):2330.
- Maroto-Morales A, Ramon M, Garcia-Alvarez O, Soler AJ, Fernandez-Santos MR, Roldan ER, Gomendio M, Perez-

<https://doi.org/10.17221/56/2025-VETMED>

- Guzman MD, Garde JJ. Morphometrically-distinct sperm subpopulations defined by a multistep statistical procedure in ram ejaculates: Intra- and interindividual variation. *Theriogenology*. 2012 May;77(8):1529-39.
- Marti JI, Aparicio IM, Garcia-Herreros M. Sperm morphometric subpopulations are differentially distributed in rams with different maturity age in cryopreserved ejaculates. *Theriogenology*. 2011 Jul 1;76(1):97-109.
- Martinez LB, Crispin RH, Mendoza MM, Gallegos OH, Martinez AA. Use of multivariate statistics to identify unreliable data obtained using CASA. *Syst Biol Reprod Med*. 2013 Jun;59(3):164-71.
- Martinez-Pastor F. What is the importance of sperm subpopulations? *Anim Reprod Sci*. 2022 Nov;246:106844.
- Martinez-Pastor F, Tizado EJ, Garde JJ, Anel L, de Paz P. Statistical series: Opportunities and challenges of sperm motility subpopulation analysis. *Theriogenology*. 2011 Mar 15;75(5):783-95.
- Menon AG, Barkema HW, Wilde R, Kastelic JP, Thundathil JC. Associations between sperm abnormalities, breed, age, and scrotal circumference in beef bulls. *Can J Vet Res*. 2011 Oct;75(4):241-7.
- Mikkola M, Desmet KJL, Kommissrud E, Riegler MA. Recent advancements to increase success in assisted reproductive technologies in cattle. *Anim Reprod*. 2024 Aug 12;21(3):e20240031.
- Muino R, Pena AI, Rodriguez A, Tamargo C, Hidalgo CO. Effects of cryopreservation on the motile sperm subpopulations in semen from Asturiana de los Valles bulls. *Theriogenology*. 2009 Oct 1;72(6):860-8.
- Muino R, Tamargo C, Hidalgo CO, Pena AI. Identification of sperm subpopulations with defined motility characteristics in ejaculates from Holstein bulls: Effects of cryopreservation and between-bull variation. *Anim Reprod Sci*. 2008 Dec;109(1-4):27-39.
- Murphy EM, Kelly AK, O'Meara C, Eivers B, Lonergan P, Fair S. Influence of bull age, ejaculate number, and season of collection on semen production and sperm motility parameters in Holstein Friesian bulls in a commercial artificial insemination centre. *J Anim Sci*. 2018 Jun;96(6):2408-18.
- Ntemka A, Tsousis G, Brozos C, Kioassis E, Boscos CM, Tsakmakidis IA. Breed differences of bull frozen-thawed semen. *Reprod Domest Anim*. 2016 Dec;51(6):945-52.
- Pena AI, Adan S, Quintela LA, Becerra JJ, Herradon PG. Relationship between motile sperm subpopulations identified in frozen-thawed dog semen samples and their ability to bind to the zona pellucida of canine oocytes. *Reprod Domest Anim*. 2018 Nov;53(Suppl\_3):14-22.
- Perez-Marin CC, Arando A, Maroto-Molina F, Marin A, Delgado JV. Sperm subpopulations and quality in fractions obtained after single layer centrifugation in fresh normospermic ram samples. *Rev Mex Cienc Pecu*. 2021 Aug;12(2):386-401.
- Ramon M, Soler AJ, Ortiz JA, Garcia-Alvarez O, Maroto-Morales A, Roldan ER, Garde JJ. Sperm population structure and male fertility: An intraspecific study of sperm design and velocity in red deer. *Biol Reprod*. 2013 Nov;89(5):110.
- Santolaria P, Vicente-Fiel S, Palacin I, Fantova E, Blasco ME, Silvestre MA, Yaniz JL. Predictive capacity of sperm quality parameters and sperm subpopulations on field fertility after artificial insemination in sheep. *Anim Reprod Sci*. 2015 Dec;163:82-8.
- Silvestre MA, Soler C, Moce E, Roldan ERS, Yaniz JL. The importance of studying factors that affect the in vitro evaluation of semen quality to predict potential fertility in males. *Biology (Basel)*. 2023 Feb 2;12(2):235.
- Solis JM, Sevilla F, Silvestre MA, Araya-Zuniga I, Roldan ERS, Saborio-Montero A, Valverde A. Effect of thawing procedure and thermo-resistance test on sperm motility and kinematics patterns in two bovine breeds. *Animals (Basel)*. 2024 Sep 25;14(19):2768.
- Souza AH, Bessoff HJ, Danzeisen E. Effect of semen type (cooled-fresh vs. frozen-thawed) on fertility of lactating dairy cows. *J Anim Sci*. 2016 Oct;94(Suppl\_5):180.
- Valverde A, Arean H, Fernandez A, Bompard D, Garcia-Molina A, Lopez-Viana J, Soler C. Combined effect of type and capture area of counting chamber and diluent on Holstein bull sperm kinematics. *Andrologia*. 2019 May;51(4):e13223.
- Valverde A, Arenan H, Sancho M, Contell J, Yaniz J, Fernandez A, Soler C. Morphometry and subpopulation structure of Holstein bull spermatozoa: Variations in ejaculates and cryopreservation straws. *Asian J Androl*. 2016 Nov-Dec;18(6):851-7.
- Viquez L, Sevilla F, Araya-Zuniga I, Soler C, Barquero V, Roldan ERS, Valverde A. Morphometric assessment of cryopreserved livestock bull spermatozoa in the tropics. *Reprod Domest Anim*. 2023 Oct;58(10):1439-47.
- Yaniz JL, Palacin I, Caycho KS, Soler C, Silvestre MA, Santolaria P. Determining the relationship between bull sperm kinematic subpopulations and fluorescence groups using an integrated sperm quality analysis technique. *Reprod Fertil Dev*. 2018 Jun;30(6):919-23.
- Yaniz JL, Vicente-Fiel S, Soler C, Recreo P, Carretero T, Bono A, Berne JM, Santolaria P. Comparison of different statistical approaches to evaluate morphometric sperm subpopulations in men. *Asian J Androl*. 2016 Nov-Dec;18(6):819-23.

Received: August 11, 2025

Accepted: November 7, 2025

Published online: January 26, 2026